

### 9.3.4 Multipoint boundary value problem with multiple solutions

Consider here a 4th-order nonlinear eigenvalue problem with multipoint boundary condition

$$u'''' = \beta z(1 + u^2), \quad u(0) = u'(1) = u''(1) = 0, \quad u'''(0) = u'''(\alpha), \quad (9.52)$$

where  $\alpha \in (0,1)$  and  $\beta$  are given constants. Writing  $u(z) = \lambda\theta(z)$  with the definition  $\lambda = u(1)$ , the original equation (9.52) becomes


$$\theta'''' = \beta z(1 + \lambda^2\theta^2), \quad \theta(0) = \theta'(1) = \theta''(1) = 0, \quad \theta'''(0) = \theta'''(\alpha), \quad (9.53)$$

with one additional boundary condition


$$\theta = \dots \quad (9.54)$$

The two solutions can be found out by means of the **BVPh 1.0**, as shown below.

**Fig. 9.21** Comparison of two eigenfunctions of (9.53) and (9.54) when  $\alpha = 1/5$  and  $\beta = 10$  by means of different initial guesses  $\lambda_0$  of the eigenvalue. Solid line: the 1st eigenfunction  $\theta(z)$  given by  $\lambda_0 = 0.63313$  and  $c_0 = -3/2$ ; Dashed line with open circles: the 2nd eigenfunction  $\theta(z)$  given by  $\lambda_0 = 2.14591$  and  $c_0 = -1/2$ .



**Fig. 9.22** Two solutions of the original equation (9.52) when  $\alpha = 1/5$  and  $\beta = 10$ . Solid line: 1st solution  $u(z)$  given by  $\lambda_0 = 0.63313$  and  $c_0 = -3/2$ ; Dashed line: 2nd solution  $u(z)$  given by  $\lambda_0 = 2.14591$  and  $c_0 = -1/2$ .



$\alpha = 0.2, \beta = 10$   
 given by  $\lambda_0 = 0.63313$  and  $c_0 = -3/2$ ; Dashed line: 2nd solution  $u(z)$  given by  $\lambda_0 = 2.14591$  and  $c_0 = -1/2$ .